

## Technical Data Sheet

### HexA-His/HexB-FLAG Heterodimer Protein

**Catalog Number:** 601001, 601002

**Size:** 25 ug, 100 ug

**Target Name:** Hexosaminidase A, Hexosaminidase B

**Regulatory Status:** RUO

#### Product Details

---

**Application:** ELISA

**Format:** Liquid, Purified

**Expression Host:** CHO

**Species:** Human

**Accession Number:** P07686/P06865

**Sources:** Recombinant human HexB (Ala43-Met556) with C-terminus DYKDDDDK tag and human HexA (Leu23-Thr529) with C-terminus His tag are co-expressed in CHO cells. Heterodimer protein is purified sequentially by anti-FLAG tag resin and Nickel resin.

**Molecular Weight:** Human HexA protein has the predicted molecular weight of 60 kD and human HexB protein has the predicted molecular weight of 59.8 kD . Under DTT-reducing conditions, both proteins migrate at approximately 65 kDa on SDS-PAGE.

**Affinity Tag:** C-His and C-DYKDDDDK

**Purity:** >95% based on SDS-PAGE under reducing condition

**Formulation:** 25 mM Tris, 150 mM NaCl, pH 7.5 (0.2 um filtered)

**Endotoxin level:** Not tested

**Protein Concentration:** 25µg size is bottled at 0.2mg/mL concentration. 100 µg size is supplied at a lot-specific concentration.

**Storage and Handling:** Briefly centrifuge the vial upon receipt. An unopened vial can be stored at 4°C for up to 2 weeks, or at -20°C or below for up to six months. The protein may be further diluted to 0.1 mg/mL using 0.22 µm-filtered 25 mM Tris, 150 mM NaCl, pH 7.5. For long-term storage, the diluted stock solution should be aliquoted and stored at  $\leq -70^{\circ}\text{C}$  to minimize freeze-thaw cycles. If additional dilution is required, carrier proteins such as FBS or BSA should be added to maintain protein stability.


#### Background Information

---

Beta-hexosaminidases are lysosomal enzymes that hydrolyze terminal N-acetyl-D-hexosamine residues from GM2 gangliosides and globo-sphingolipids. They exist in three isoforms: Hex A ( $\alpha\beta$ ), Hex B ( $\beta\beta$ ), and Hex S ( $\alpha\alpha$ ), formed by different combinations of  $\alpha$  and  $\beta$  subunits encoded by the HEXA and HEXB genes. Recombinant HexA-His/HexB-FLAG Heterodimer corresponds to Hex A isoform.

#### Product Data

---



Purified HexA/B heterodimer (HexA-His/HexB-FLAG) and final products on SDS-PAGE under non-reducing (P-) and reducing (P+) conditions. The purity of HexA/B heterodimer is greater than 95%.