

## Technical Data Sheet

### Biotin Anti-Human CD197

**Catalog Number:** 107307, 107308

**Size:** 25 ug, 100 ug

**Target Name:** CD197, CCR7, BLR2, CDw197, EBI1, CMKBR7

**Regulatory Status:** RUO

#### Product Details

---

**Clone:** 197AM2a

**Application:** FC

**Reactivity:** Human

**Format:** Biotin

**Isotype:** Mouse IgG1

**Antibody Type:** Monoclonal

**Formulation:** Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide

**Protein Concentration:** 0.2 mg/mL

**Storage and Handling:** The antibody solution should be stored between 2°C and 8°C

**Recommended Usage:** For flow cytometric staining, it is recommended to use less than 0.25 ug of this reagent per 0.5-1.0 million cells in a 100 µL volume. Optimal reagent performance should be determined by titration for each specific application


#### Background Information

---

C-C chemokine receptor type 7 is a protein that in humans is encoded by the CCR7 gene. CD197 is a chemokine receptor that binds CCL19 and CCL21. CCR7 is expressed on naive T cells, regulatory T cell, memory T cell, dendritic cell, mature B cell. Through regulating the homing of CD197 expressing cells into secondary lymphoid organs, it affects the interaction among these cells and thus the immune response.

#### Product Data

---



Human peripheral blood lymphocytes stained with biotinylated Anti-Human CD197 clone 197AM2a (blue histogram) or an isotype control (gray histogram) , followed by SA-PE.