

InnoCyto Inc.

15375 Barranca Pkwy, Suite I-103 Irvine, CA 92618

Technical Data Sheet

APC Anti-Human CD14

Catalog Number: 110107, 110108

Size: 25 tests, 100 tests

Target Name: CD14, LPS receptor

Regulatory Status: RUO

Product Details

Clone: M5E2
Application: FC
Reactivity: Human
Format: APC

Isotype: Mouse IgG2a

Antibody Type: Monoclonal

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v)

BSA

Protein Concentration: Supplied at a lot-specific concentration.

Storage&Handling: The antibody solution should be stored undiluted between 2°C and 8°C, and

protected from prolonged exposure to light. Do not freeze.

Recommended Usage: For flow cytometric staining, it is recommended to use 5 uL of this reagent per 0.5-1.0 million cells in a 100 μ L volume. Optimal reagent performance should be determined by

titration for each specific application. **Excitation Laser:** Red Laser (633 nm)

Release Date: Dec-25

Background Information


CD14 is a pattern recognition receptor critical for innate immunity. It exists as a glycosylphosphatidylinositol (GPI)-anchored membrane protein (mCD14) on monocytes, macrophages, and neutrophils, and as a soluble protein (sCD14) in the blood, generated via enzymatic cleavage or shedding. Membrane CD14 acts as a pro-inflammatory co-receptor for lipopolysaccharide (LPS), interacting with LPS-binding protein (LBP) and TLR4 to activate NF-?B and MAPK signaling, driving inflammatory responses. Soluble CD14 enables LPS responsiveness in cells lacking CD14, such as endothelial cells. CD14 expression is modulated by stimuli including LPS, IFN?, PMA, calcium ionophore, and anti-CD14 antibodies. Elevated sCD14 levels are observed in conditions such as sepsis, systemic lupus erythematosus, HIV infection, common variable immunodeficiency (CVID), nonalcoholic fatty liver disease (NAFLD), and breast invasive ductal carcinoma, highlighting its potential as a biomarker for infection, inflammation, and disease prognosis.

Product Data



InnoCyto Inc.

15375 Barranca Pkwy, Suite I-103 Irvine, CA 92618



Human peripheral blood monocytes was stained with APC Anti-Human CD14 clone M5E2 (color-filled histogram) or an isotype control (gray histogram).