

## In Vivo Star Anti-Mouse CD25 (IL-2R $\alpha$ ) Antibody

|                           |                                                                        |
|---------------------------|------------------------------------------------------------------------|
| <b>Catalog Number:</b>    | 508901, 508902, 508903                                                 |
| <b>Size:</b>              | 1 mg, 5 mg, 25 mg                                                      |
| <b>Target Name:</b>       | IL-2 receptor $\alpha$ chain, Low affinity IL-2R, IL-2R $\alpha$ chain |
| <b>Regulatory Status:</b> | RUO                                                                    |

### PRODUCT DETAILS

---

|                             |                                                                                                                         |
|-----------------------------|-------------------------------------------------------------------------------------------------------------------------|
| <b>Clone:</b>               | PC61.5.3-m2c                                                                                                            |
| <b>Application:</b>         | ELISA, WB, Flow cytometry, IHC, ICC, animal model study                                                                 |
| <b>Reactivity:</b>          | Mouse                                                                                                                   |
| <b>Format:</b>              | Liquid                                                                                                                  |
| <b>Product Description:</b> | In vivo Grade Recombinant Anti-mouse CD25 Monoclonal Antibody                                                           |
| <b>Isotype:</b>             | Mouse IgG2c Kappa                                                                                                       |
| <b>Antibody Type:</b>       | Recombinant                                                                                                             |
| <b>Purity:</b>              | >95% by reducing SDS-PAGE                                                                                               |
| <b>Endotoxin:</b>           | < 1 EU per 1 mg of the protein by the LAL method.                                                                       |
| <b>Storage Conditions:</b>  | 4°C                                                                                                                     |
| <b>Grade:</b>               | In vivo                                                                                                                 |
| <b>Recommended Usage:</b>   | This product is suitable for in vivo animal use. Optimal amounts need to be determined empirically for each experiment. |
| <b>Hidden Synonyms:</b>     | InVivoMab, InVivoPlus, GoInVivo, In Vivo Gold                                                                           |
| <b>RRID:</b>                | AB_3739363                                                                                                              |

### BACKGROUND INFORMATION

---

CD25, also known as the interleukin-2 receptor alpha chain (IL-2R $\alpha$ ), is a transmembrane glycoprotein that plays a central role in regulating immune responses. It functions as part of the interleukin-2 (IL-2) receptor complex, which is essential for T cell proliferation, survival, and differentiation. CD25 itself has low affinity for IL-2 when expressed alone, but when combined with IL-2 receptor beta (CD122) and the common gamma chain (CD132), it forms the high-affinity IL-2 receptor complex capable of effective signal transduction.

Structurally, CD25 is a single-pass type I membrane protein composed of an extracellular domain of approximately 219 amino acids responsible for IL-2 binding, a hydrophobic transmembrane segment, and a short cytoplasmic tail that lacks intrinsic signaling domains. The extracellular region is heavily glycosylated, which stabilizes its conformation and facilitates ligand interaction. Because the alpha chain alone is not signaling-competent, it acts primarily to increase the receptor complex's affinity for IL-2 and to expand the range of cells responsive to low cytokine concentrations.

CD25's main ligand, IL-2, is a cytokine crucial for T lymphocyte expansion and immune tolerance. Engagement of IL-2 with the high-affinity receptor triggers the JAK-STAT signaling pathway, leading to cell proliferation, differentiation, and regulatory T cell (Treg) function. CD25 is constitutively expressed on Tregs and upregulated on activated CD4+ and CD8+ T cells, making it a marker of immune activation as well as immune regulation.

Aberrant CD25 expression or IL-2 signaling contributes to immune dysregulation and disease. In autoimmune disorders such as multiple sclerosis and type 1 diabetes, alterations in the IL-2/CD25 axis impair Treg function and tolerance mechanisms. Elevated CD25 expression is also found in certain malignancies, particularly adult T-cell leukemia/lymphoma and Hodgkin lymphoma, where it may serve as a biomarker of malignant proliferation. Moreover, soluble CD25, released from cell surfaces, can act as a decoy receptor, modulating IL-2 availability and contributing to immune suppression in cancer and chronic inflammation.

Therapeutically, CD25 is a prominent target for immune modulation. Monoclonal antibodies such as basiliximab and daclizumab have been developed to block IL-2 binding, preventing T cell activation and mitigating graft rejection in organ transplantation. Conversely, IL-2 or CD25-targeted therapies that enhance regulatory T cell function are being explored to treat autoimmune diseases and promote immune tolerance. Thus, CD25 remains a critical immunological node, balancing activation and regulation within the immune system.

This product is supplied subject to the terms and conditions at [www.innocyto.com/web/terms.php](http://www.innocyto.com/web/terms.php) and may only be used as provided in the stated terms. Products are for Research Use Only.