

## Mouse Fc blocking reagent

<b>Catalog Number:</b>	201813, 201814
<b>Size:</b>	100 &mu;g, 500 &mu;g
<b>Target Name:</b>	CD16/32
<b>Regulatory Status:</b>	RUO

### PRODUCT DETAILS

---

<b>Clone:</b>	2.4G2
<b>Application:</b>	Flow Cytometry
<b>Reactivity:</b>	Mouse
<b>Format:</b>	Purified
<b>Isotype:</b>	Rat IgG2b
<b>Antibody Type:</b>	Monoclonal
<b>Formulation:</b>	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide
<b>Protein Concentration:</b>	0.5 mg/mL
<b>Storage&amp;Handling:</b>	The antibody solution should be stored between 2°C and 8°C
<b>Recommended Usage:</b>	For flow cytometric analysis, block Fc receptors by pre-incubating cells with a mouse Fc blocker for 5-10 minutes on ice at 1.0 µg per 10 <sup>6</sup> cells in a volume of 100 µL prior to immunostaining. Washing between the blocking step and primary antibody staining is not required.
<b>Isotype Controls:</b>	N/A
<b>Antibody Family:</b>	Mouse Antibodies


### BACKGROUND INFORMATION

---

CD16/32 are Fc-gamma receptors (FcγRs) expressed on a variety of immune cells, including B cells, monocytes/macrophages, NK cells, granulocytes, mast cells, and dendritic cells. CD16 corresponds to the low-affinity Fc receptor III (FcγRIII), while CD32 corresponds to Fc receptor II (FcγRII). These receptors bind antibody-antigen immune complexes, linking innate and adaptive immunity and mediating adaptive immune responses. In research, antibodies against CD16/CD32 are commonly used to block Fc receptor-mediated interactions, preventing non-specific binding of antibodies or immunoglobulin complexes to immune cells during experiments such as flow cytometry and immunohistochemistry, thereby improving experimental accuracy.

PRODUCT DATA

---



Mouse splenocytes were incubated with Innocyto Mouse Fc blocking reagent (1 ug/1e6 cells; color-filled histogram) and then stained with FITC anti-Mouse CD90. 2 clone 30-H12. Untreated cells are shown as the dotted-line histogram.

This product is supplied subject to the terms and conditions at [www.innocyto.com/web/terms.php](http://www.innocyto.com/web/terms.php) and may only be used as provided in the stated terms. Products are for Research Use Only.