

Anti-Human HLA-DR/DP/DQ Antibody

Catalog Number:	118901, 118902
Size:	25 µg, 100 µg
Target Name:	HLA-DR, HLA-DP, and HLA-DQ, MHC class II, pan HLA-II, pan MHC-II
Regulatory Status:	RUO

PRODUCT DETAILS

Clone:	IVA12
Application:	Flow Cytometry
Reactivity:	Human
Format:	Purified
Isotype:	Mouse IgG1
Antibody Type:	Monoclonal
Formulation:	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide
Protein Concentration:	0.5 mg/mL
Storage&Handling:	The antibody solution should be stored between 2°C and 8°C
Isotype Control:	301401

BACKGROUND INFORMATION

Human HLA-DR, HLA-DP, and HLA-DQ are the three major classical human leukocyte antigen (HLA) class II molecules encoded within the major histocompatibility complex (MHC) on chromosome 6. These cell-surface glycoproteins are primarily expressed on professional antigen-presenting cells, including dendritic cells, macrophages, and B lymphocytes. Their principal function is to present peptides derived from extracellular proteins to CD4+ T helper cells, initiating and regulating adaptive immune responses. This process is essential for host defense against pathogens and for the development of immune memory.


Each HLA class II molecule consists of a polymorphic α (alpha) chain and β (beta) chain that together form a peptide-binding groove capable of accommodating peptides approximately 13–25 amino acids in length. The groove binds peptides generated in endosomal and lysosomal compartments following antigen processing. Peptide loading is facilitated by accessory molecules such as HLA-DM after removal of the invariant chain-derived CLIP peptide. The extensive genetic polymorphism of HLA-DR, HLA-DP, and HLA-DQ enables presentation of a broad repertoire of peptide ligands, including self and foreign antigens.

Specific HLA class II alleles are strongly associated with autoimmune diseases. For example, HLA-DR and HLA-DQ variants contribute to susceptibility to type 1 diabetes, rheumatoid arthritis, celiac disease, multiple sclerosis, and systemic lupus erythematosus. Certain HLA-DP alleles have also been linked to chronic hepatitis B outcomes and other immune-mediated disorders. These molecules are important in transplantation because mismatched HLA alleles increase the risk of graft rejection.

Therapeutically, HLA-DR, HLA-DP, and HLA-DQ are used in tissue matching for organ and stem cell transplantation, disease risk assessment, vaccine development, and immunotherapy. Their ability to present antigenic peptides also makes them attractive

targets for antigen-specific tolerance strategies and personalized immunotherapies designed to modulate pathogenic T-cell responses while preserving protective immunity.

PRODUCT DATA



Human peripheral blood lymphocytes were stained with i F647 Anti-Human CD19 clone Cyte A-SJ25C1 and Purified Anti-Human HLA-DR/DP/DQ clone IVA12 (left) or an isotype control (right), followed by PE anti-Mouse IgG.

This product is supplied subject to the terms and conditions at www.innocyto.com/web/terms.php and may only be used as provided in the stated terms. Products are for Research Use Only.