

InnoCyto Inc.

15375 Barranca Pkwy, Suite I-103 Irvine, CA 92618

Technical Data Sheet

Anti-Mouse/human CD45R/B220

Catalog Number: 200901, 200902

Size: 100 ug, 500 ug

Target Name: CD45R, B220 Regulatory Status: RUO

Product Details

Clone: RA3-6B2 Application: FC

Reactivity: Human, Mouse

Format: Purified Isotype: Rat IgG2a

Antibody Type: Monoclonal

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide

Protein Concentration: 0.5 mg/mL

Storage&Handling: The antibody solution should be stored between 2°C and 8°C

Recommended Usage: For flow cytometric staining, it is recommended to use less than 0.2 ug of this reagent per 0.5-1.0 million cells in a 100 μ L volume. Optimal reagent performance should be

determined by titration for each specific application

Release Date: Aug-25

Background Information


CD45R, also known as B220, is an isoform of CD45 (PTPRC), a member of the protein tyrosine phosphatase (PTP) family. With a molecular weight of approximately 180–240 kDa, CD45R is a transmembrane glycoprotein expressed on B cells at all developmental stages, as well as on activated B cells, subsets of T and NK cells, and certain abnormal T cell populations involved in systemic autoimmunity in MRL-Faslpr and MRL-Fasgld mice. Structurally, CD45R consists of an extracellular domain, a single transmembrane region, and two tandem intracytoplasmic catalytic domains. Functionally, CD45R plays a critical role in T cell receptor (TCR) and B cell receptor (BCR) signaling by regulating Src-family kinases, which are essential for lymphocyte activation and development. The primary ligands for CD45 include galectin-1, CD2, CD3, and CD4. CD45R is widely used as a pan-B cell marker, though CD19 may provide greater specificity for B cells.

Product Data



InnoCyto Inc.

15375 Barranca Pkwy, Suite I-103 Irvine, CA 92618



Mouse splenocytes were stained with purified Anti-Human/Mouse CD45R/B220 clone RA3-6B2 (color-filled histogram) or an isotype control (gray histogram), followed by PE anti-rat IgG 2nd antibody.