www.innocyto.com

InnoCyto Inc.

15375 Barranca Pkwy, Suite I-103
Irvine, CA 92618
Email: info@innocyto.com
Phone: 1-949-418-7543

PE Anti-Human CD274 (PD-L1) Antibody

Catalog Number:	120005, 120006
Size:	25 tests, 100 tests
Target Name:	CD274, PD-L1, B7-H1, Programmed cell death ligand 1, B7 homolog 1
Regulatory Status:	RUO

PRODUCT DETAILS

Clone:	29E.2A3
Reactivity:	Human
Format:	PE
Isotype:	Mouse IgG2b
Antibody Type:	Monoclonal
Formulation:	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA
Protein Concentration:	Supplied at a lot-specific concentration.
Storage&Handling:	The antibody solution should be stored undiluted between 2°C and 8°C, and protected from prolonged exposure to light. Do not freeze.
Recommended Usage:	For flow cytometric staining, it is recommended to use 5 µL of this reagent per 0.5-1.0 million cells in a 100 µL volume. Optimal reagent performance should be determined by titration for each specific application. PE has an excitation max at 565 nm and an emission max at 575 nm.
Excitation Laser:	Blue Laser (488 nm) Green/Yellow laser (532/561nm)
Isotype Controls:	301615
Antibody Family:	Human Antibodies
Storage & Handling:	The antibody solution should be stored undiluted between 2°C and 8°C, and protected from prolonged exposure to light. Do not freeze.
Isotype Control:	301615

BACKGROUND INFORMATION

Human programmed death-ligand 1 (PD-L1), also known as CD274 or B7-H1, is an immune checkpoint protein that plays a central role in regulating T-cell activity and maintaining peripheral immune tolerance. PD-L1 is expressed on antigen-presenting cells, including dendritic cells and macrophages, and can be induced on many other cell types by inflammatory cytokines such as interferon-γ. It is frequently upregulated by tumor cells as a mechanism of immune evasion.

PD-L1 is a type I transmembrane glycoprotein of approximately 290 amino acids. Its extracellular region contains an N-terminal immunoglobulin variable (IgV)-like domain and a membrane-proximal IgC2-like domain, followed by a single transmembrane helix and a short cytoplasmic tail of approximately 30 amino acids. The IgV domain contains the primary binding interface for PD-1.

Structural studies show that PD-L1 engages PD-1 through an extensive protein-protein interface involving hydrophobic and polar interactions.

The principal receptor for PD-L1 is programmed cell death protein 1 (PD-1/CD279), expressed mainly on activated T cells and other immune cells. PD-L1 can also interact with B7-1 (CD80), including interactions occurring in cis on antigen-presenting cells. Engagement of PD-1 by PD-L1 suppresses T-cell receptor signaling, proliferation, cytokine production, and cytotoxic activity, thereby limiting excessive immune responses.

Dysregulated PD-L1 expression contributes to immune suppression in cancer and has been associated with chronic infections and inflammatory diseases. Therapeutically, the PD-1/PD-L1 pathway is one of the most important targets in cancer immunotherapy. Monoclonal antibodies blocking PD-1 or PD-L1 can restore antitumor T-cell activity and have demonstrated substantial clinical benefit across multiple cancers. PD-L1 is therefore both a therapeutic target and an important biomarker for patient selection in certain immunotherapy settings.

PRODUCT DATA



Human peripheral blood mononuclear cells stimulated with Anti-Human CD3/CD28 for 3-days were stained with PE Anti-Human CD274 (PD-L1) clone 29E.2A3 (color-filled histogram) or an isotype control (gray histogram).

This product is supplied subject to the terms and conditions at www.innocyto.com/web/terms.php and may only be used as provided in the stated terms. Products are for Research Use Only.