

Anti-Human B7-H3 (Ifinatumab Biosimilar)

Catalog Number:	503201, 503202, 503203
Size:	1 mg, 5 mg, 20 mg
Target Name:	CD276, B7-H3, B7H3, B7RP-2
Regulatory Status:	RUO

PRODUCT DETAILS

Clone:	Ifinatumab
Application:	Flow cytometry, animal model study
Reactivity:	Human
Format:	Liquid
Product Description:	Ifinatumab Biosimilar, Human B7-H3 Monoclonal Antibody
Isotype:	Human IgG1
Clonality:	Recombinant
Immunogen:	Human B7-H3
Species specificity:	Human
Purity:	>95% by reducing SDS-PAGE
Grade:	In vivo
Min Sample Size:	1 mg
Storage Conditions:	4°C
Maximal Shelf Life:	12 months
Synonyms:	CD276
Antibody Type:	Recombinant
Reactivity:	Human

BACKGROUND INFORMATION

Ifanatumab, also known as Ifinatumab deruxtecan or DS-7300a, is a humanized monoclonal antibody belonging to the immunoglobulin G1 kappa (IgG1κ) subclass. It is engineered to recognize and bind to the human B7-H3 (CD276) cell surface glycoprotein, a member of the B7 family involved in immune regulation and intercellular signaling. As a therapeutic, the Ifanatumab molecule is part of an antibody-drug conjugate (ADC) system, where the IgG1 antibody is covalently linked to a cytotoxic topoisomerase I inhibitor, deruxtecan, via a cleavable peptide-based linker. The complete conjugate typically displays a molecular weight near 150 kilodaltons (kDa) and maintains the classical Y-shaped antibody configuration.

The antibody portion of Ifanatumab consists of two heavy chains and two light chains joined by disulfide bonds. The variable (VH

and VL) domains form the antigen-binding regions, containing complementarity-determining regions (CDRs) that confer high specificity toward epitopes on the extracellular domains of B7-H3. The binding mechanism involves non-covalent interactions, including hydrogen bonds and hydrophobic contacts, ensuring nanomolar-level affinity. Upon binding to B7-H3 on the cell surface, the ADC-antigen complex undergoes receptor-mediated endocytosis, transporting the conjugate into the lysosomal compartment of the target cell.

The Fc (fragment crystallizable) region of the IgG1 backbone supports molecular stability and prolonged half-life by engaging neonatal Fc receptors (FcRn), while maintaining minimal immune effector activation.